


Asymmetric hydrogenation of olefins

- Monodentate phosphoramidites. Affordable ligands for asymmetric hydrogenation
- MonoPhos™ one of the most effective ligands is synthesised in one step!
- The rhodium catalysed olefin hydrogenation using monodentate phosphoramidites as ligands has been used to synthesise α - and β -amino acids with very high enantioslectivity.

Asymmetric hydrogenation of olefins

- Synthesis of ligands



References

1. M. van den Berg, A. J. Minnaard, E. P. Schudde, J. van Esch, A. H. M. de Vries, J. G. de Vries, B. L. Feringa, *J. Am. Chem. Soc.* 2000, 122, 11539.
2. M. van den Berg, A. J. Minnaard, J. G. de Vries, B. L. Feringa (DSM N.V.), World Patent WO 02/04466, 2002.
3. M. van den Berg, A. J. Minnaard, R.M. Haak, M. Leeman, E.P. Schudde, A. Meetsma, B.L. Feringa, A.H.M. de Vries, C.E.P. Maljaars, C.E. Willans, D. Hyett, J.A.F. Boogers, H. J.W. Henderickx and J.G. de Vries, *Adv. Synth. Catal.*, 2003,.(Noyori issue) accepted for publication.
4. D. Peña, A.J. Minnaard, J.G. de Vries and B.L. Feringa, *J. Am. Chem. Soc.*, accepted for publication